

Holbrook Primary School

Individual protocol for an EpiPen adrenaline auto injector

Child's name:

D.O.B.

Class:

Nature of Allergy:

Contact Information

Name					Relationship to pupil			
Phone numbers	Work		Home		Mobile		Other	

If I am unavailable, please contact:

Name					Relationship to pupil			
Phone numbers	Work		Home		Mobile		Other	

GP

Name:

Phone No:

Address:

Clinic/ Hospital Contact

Name:

Phone No:

Address:

MEDICATION EPIPEN

Name on EPIPEN 1 & Expiry date:

Name on EPIPEN 2 & Expiry date:

- It is the parents responsibility to supply 2 EPIPEN auto injectors and to ensure they have not expired

Dosage & Method: **1 DOSE INTO UPPER OUTER THIGH**

- The school staff will take all reasonable steps to ensure does not eat any food items unless they have been prepared / approved by parents
- It is the schools responsibility to ensure this care plan is reviewed and parents inform the school of any changes in condition or treatment.

Agreed by: School Representative.....Date.....

- I agree that the medical information contained in this plan may be shared with individuals involved with my child's care and education.

TO ACCOMPANY HEALTHCARE PLAN

- I give my consent for the school to administer my child's Epipen **or the school held adrenaline auto-injector** (if my child's pen is lost/forgotten or malfunctions) to be administered in an emergency as detailed in this plan

Signed:.....Print name..... Date.....

I am the person with parental responsibility

TO ACCOMPANY HEALTHCARE PLAN

Individual protocol for using an Epipen (Adrenaline Auto injector)

PLEASE AMEND THE SYPTOMS TO REFLECT YOUR CHILD IF KNOWN.

Symptoms may include:

- Difficulty in swallowing / speaking / breathing
- Wheezy / irregular breathing / excessive coughing
- Hoarseness
- Nettle rash (hives) anywhere on body
- Sense of impending doom
- Swelling of throat and mouth
- Abdominal pain, nausea & vomiting
- Feeling of weakness (BP drops)
- Collapse & unconsciousness
- Cold and clammy

Stay Calm

Reassure

**One member of staff
to Dial 999**

REMEMBER

**A = AIRWAY
B = BREATHING
C = CIRCULATION**

Give EPIPEN first then dial 999

Administer Epipen in the upper outer thigh

Remove grey safety cap
Hold epipen with black tip
downwards against thigh
jab firmly.

Hold epipen in place for 10 seconds

Can be given through clothing,
but not very thick clothing.
Note time of injection given

**If no improvement give
2nd EPIPEN 5 minutes
later**

Call Parents

Reassure

.....

Telephoning for an ambulance

You need to say: "I have a child in anaphylactic shock".

Give school details: Holbrook Primary School, Holbrook School Lane, Horsham, RH12 5PP

Give details: Childs name has a severe allergy and what has happened.

**DO NOT PUT THE PHONE DOWN UNTIL YOU ARE SURE ALL THE NECESSARY INFORMATION HAS
BEEN GIVEN**

Someone to wait by the school gate to direct the ambulance staff straight to the child.